

obtained directly from total DNA or from total RNA. The cDNA clones are not naturally occurring, but rather are preferably obtained via manipulation of a partially purified naturally occurring substance (messenger RNA). The construction of a cDNA library from mRNA involves the creation of a synthetic substance (cDNA) and pure individual cDNA clones can be isolated from the synthetic library by clonal selection of the cells carrying the cDNA library. Thus, the process which includes the construction of a cDNA library from mRNA and isolation of distinct cDNA clones yields an approximately 10^6 -fold purification of the native message. Thus, purification of at least one order of magnitude, preferably two or three orders, and more preferably four or five orders of magnitude is expressly contemplated.

By "enriched" in reference to nucleic acid is meant that the specific DNA or RNA sequence constitutes a significantly higher fraction (2 - 5 fold) of the total DNA or RNA present in the cells or solution of interest than in normal or diseased cells or in the cells from which the sequence was taken. This could be caused by a person by preferential reduction in the amount of other DNA or RNA present, or by a preferential increase

in the amount of the specific DNA or RNA sequence, or by a combination of the two. However, it should be noted that enriched does not imply that there are no other DNA or RNA sequences present, just that the relative amount of the sequence of interest has been significantly increased in a useful manner and preferably separate from a sequence library. The term "significantly" here is used to indicate that the level of increase is useful to the person making such an increase, and generally means an increase relative to other nucleic acids of about at least 2 fold, more preferably at least 5 to 10 fold or even more. The term also does not imply that there is no DNA or RNA from other sources. The DNA from other sources may, for example, comprise DNA from a yeast or bacterial genome, or a cloning vector such as pUC19. This term distinguishes from naturally occurring events, such as viral infection, or tumor type growths, in which the level of one mRNA may be naturally increased relative to other species of mRNA. That is, the term is meant to cover only those situations in which a person has intervened to elevate the proportion of the desired nucleic acid.

By "recombinant" in reference to a nucleic acid is

meant the nucleic acid is produced by recombinant DNA techniques such that it is distinct from a naturally occurring nucleic acid.

By "comprising" is meant including, but not limited to, whatever follows the word "comprising". Thus, use of the term

5 "comprising" indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be present. By "consisting of" is meant including, and limited to, whatever follows the phrase "consisting of". Thus, the phrase "consisting of" indicates that the listed elements are required or mandatory, and that no other elements may be present.

By "consisting essentially of" is meant including any elements listed after the phrase, and limited to other elements that do not interfere with or contribute to the activity or action specified in the disclosure for the listed elements. Thus, the

15 phrase "consisting essentially of" indicates that the listed elements are required or mandatory, but that other elements are optional and may or may not be present depending upon whether or not they affect the activity or action of the listed elements.

The invention also features a nucleic acid probe for
20 the detection of a nucleic acid encoding the antithrombosis

enzyme, its A chain or B chain, or fragments thereof. In an example, the nucleic acid probe contains nucleic acid that will hybridize to SEQ ID NO: 1, but not to the nucleic acid sequence of the IX/X-binding protein (Matsuzaki et al., Biochem. Biophys.

5 Res. Com. 220(2):382-387, 1996) under high stringency hybridization conditions. By "high stringency hybridization conditions" is meant those hybridizing conditions that (1) employ low ionic strength and high temperature for washing, for example, 0.015 M NaCl/0.0015 M sodium citrate/0.1% SDS at 50°C; (2) employ during hybridization a denaturing agent such as formamide, for example, 50% (vol/vol) formamide with 0.1% bovine serum albumin/0.1% Ficoll/0.1% polyvinylpyrrolidone/50 mM sodium phosphate buffer at pH 6.5 with 750 mM NaCl, 75 mM sodium citrate at 42°C; or (3) employ 50% formamide, 5 x SSC (0.75 M NaCl, 15 0.075 M Sodium pyrophosphate, 5 x Denhardt's solution, sonicated salmon sperm DNA (50 g/ml), 0.1% SDS, and 10% dextran sulfate at 42°C, with washes at 42°C in 0.2 x SSC and 0.1% SDS. Under stringent hybridization conditions only highly complementary nucleic acid sequences hybridize. Preferably, such conditions 20 prevent hybridization of nucleic acids having 1 or 2 mismatches

out of 20 contiguous nucleotides.

In another aspect, the invention describes a recombinant cell or tissue containing an exogenous nucleic acid coding for the antithrombosis enzyme, its A chain or B chain, or fragments thereof.

In another aspect, this invention features a pharmaceutical composition for reducing or eliminating thrombosis in a human or animal subject. This composition contains a pharmaceutically effective amount of the antithrombosis enzyme and a pharmaceutically acceptable carrier.

By "pharmaceutically effective amount" is meant an amount of a pharmaceutical compound or composition having a therapeutically relevant effect on thrombosis or diseases or pathological conditions related to thrombosis. A therapeutically relevant effect prevents or relieves to some extent one or more symptoms of thrombosis or diseases or pathological symptoms related to thrombosis in a patient or returns to normal either partially or completely one or more physiological or biochemical parameters associated with or causative of thrombosis or diseases or pathological symptoms related to thrombosis, e.g., reducing or

inhibiting thrombosis; curing, reducing, or inhibiting diseases and processes that are mediated by thrombosis.

In another aspect, this invention features a method of reducing or inhibiting thrombosis in a human or animal subject by administering to the subject a pharmaceutically effective amount of the antithrombosis enzyme.

In another aspect, this invention features a method of isolating and/or purifying the antithrombosis enzyme. In this method, a crude extract containing the antithrombosis enzyme, e.g., a crude extract of the snake venom of Southern-Anhui *Agkistrodon acutus*, is dissolved in a buffer and brought into contact with an anion exchange column, the elution sample with fibrinolytic activity and antiplatelet aggregation activity is collected, concentrated and separated by gel permeation chromatography, and the elution sample with fibrinolytic activity and antiplatelet aggregation activity is collected and desalted. This invention features an antithrombosis enzyme isolated by this process.

The present invention also provides recombinant DNA molecules characterized by a DNA sequence encoding the

antithrombosis enzyme alone or fused to a DNA sequence which codes for a conventional anti-thrombosis polypeptide. The synthesis of these DNA molecules may be achieved by methods well known in the art. For example, these recombinant DNA molecules may be isolated from an *Agkistrodon acutus* venom gland cDNA library. The synthesis of cDNA libraries and the choice of vector into which the cDNA molecules may be cloned are conventional techniques. The invention also relates to hosts transformed with these recombinant DNA molecules, as well as to the recombinant products expressed by these hosts. And the present invention relates to chemically synthesized antithrombosis enzyme. Such synthetic polypeptides may be prepared by conventional chemical synthesis techniques, for example, synthesis on a solid support.

This invention further relates to pharmaceutically acceptable compositions and combinations, and methods utilizing these natural, recombinant or synthetic antiplatelet polypeptides in the treatment extracorporeal blood.

By "extracorporeal blood" is meant blood that is removed from a patient, subjected to extracorporeal treatment,

and returned to the patient in processes such as dialysis, blood
filtration or blood bypass during surgery. The term also
includes blood products which are stored extracorporeally for
eventual administration to a patient. Such products include
5 whole blood, platelet concentrates and other blood fractions in
which inhibition of platelet aggregation and platelet release is
desired.

Other features and advantages of the invention will be
apparent from the following drawing and detailed description of
the invention and from the claims.

BRIEF DESCRIPTION OF THE DRAWING

Figure 1 shows the putative cDNA sequence and amino
acid sequence of the antithrombosis enzyme, B chain. A segment
15 of about 10-20 nucleotides in the cDNA clone is being sequenced.
A leader peptide -- M G R F I F V S F G L L V V F
L S L S G T A A -- is cleaved before the assembly of A
chain and B chain to form the antithrombosis enzyme.

DETAILED DESCRIPTION OF THE INVENTION

The present invention includes compositions and methods for the treatment of diseases and pathological conditions. These diseases or pathological conditions are mediated by or associated with thrombosis. The compositions of this invention contain an antithrombosis enzyme, which can be isolated from the venom of Southern-Anhui *Agkistrodon acutus*, or synthesized by chemical or biological methods (e.g., cell culture, recombinant gene expression, or peptide synthesis). Recombinant techniques include gene amplification from DNA sources using the polymerase chain reaction (PCR), RNA polymerase based amplification, and gene amplification from RNA sources using reverse transcriptase/PCR.

Purification of the Antithrombosis enzyme from Snake Venom

The antithrombosis enzyme of this invention was extracted and purified from the venom of Southern-Anhui *Agkistrodon acutus* in China:

(1) Dissolve the crude snake venom in buffer, proceed with centrifugal precipitation and chromatographic separation through an anion exchange column, and collect roughly-isolated

fibrinolytic component by elution. The elution is conducted with both a pH linear gradient and a salt linear gradient. The pH linear gradient is from pH 7.5-8.5 to pH 6.0-7.0; and the salt linear gradient is from 0 to 0.5 M. The elution speed is 25-80 ml/hr.

(2) Concentrate this fibrinolytic component and proceed with gel permeation chromatography to get the refined antithrombosis enzyme solution.

(3) Desalt the refined antithrombosis enzyme solution and get the purified antithrombosis enzyme solution.

The above procedure was conducted at about 4-10°C.

The buffer used to dissolve the crude snake venom and the roughly-isolated elute is a Tris-HCl buffer (e.g. 0.02 M, pH 7.5-8.5 N-(tris-hydroxymethyl)-amino methane-Hydrochloric acid), or a phosphate buffer (e.g. 0.02 M, pH 7.5-8.5 $\text{Na}_2\text{HPO}_4\text{-NaH}_2\text{PO}_4$).

The anion exchange column contains weak acidic anion exchanger, e.g., DEAE Sephadex A50, or DEAE Sepharose Fast Flow.

The gel permeation chromatography uses a gel for separating substances in the range between about 5 kD and 300 kD, e.g., Sephacryl S-100HR, S-200HR, S-300HR, or Sephadex G75. The

buffer used in the gel permeation chromatography is a 0.05-0.30 M sodium chloride (NaCl) solution.

The fibrinolytic component was concentrated by ultrafiltration, blow-drying with cold air, or freeze-drying.

5 The desalting was conducted against distilled water for 1-6 hrs.

The above method for isolating and purifying the antithrombosis enzyme from the venom of *Agkistrodon acutus* is simple, provides high-purity product and shortens the time by processing the pH linear gradient elution and the salt linear gradient elution simultaneously. The shortened purification cycle increases the efficiency of industrial production.

The purification methods of this invention are also useful to isolate proteolytic fragments of the antithrombosis enzyme which retain the biological activity of the intact polypeptide. Such proteolytic fragments may occur naturally or artifactually as a result of the purification process.

The purification procedure is further described by the following three examples:

Example 1

(1) Dissolve 3 g Southern-Anhui *Agkistrodon actus* venom in 20 ml 0.02 M pH 8.0 Tris-HCl buffer. Centrifuge 15 min at 3000 rpm. Collect the supernatant. Resuspend the pellet in 5 ml above buffer and agitate well. Repeat the centrifugation once. Collect the supernatant with the earlier collection. Store at 4°C.

(2) Process the DEAE Sephadex A-50 anion exchanger with alkaline-acid and wash with distilled water to neutrality. Then wash with 0.02 M pH 8.0 Tris-HCl buffer till the eluate's pH reaches 8.0. Keep in boiling-water bath for 2 hrs to remove air bubbles and ensure swelling.

(3) Pack the column with the processed gel from Step (2) when it is cooled to ambient temperature. The column volume is 3x80 cm³. After equilibration, load the supernatant sample from Step (1) and elute with linear gradient buffer. The buffer in the stock bottle is 0.02 M Tris-HCl, pH 6.3, containing 0.5 M NaCl. The buffer in the gradient bottle is 0.02 M Tris-HCl buffer, pH 8.0. The flow rate is 42 ml/hr. Examine the eluate under UV 280 nm. Collect the eluate 48 hours after the sample is

loaded by automatic fraction collector at the rate of 7 ml per tube and 6 tubes per hour. 12 elution peaks are obtained, among which the 6th is the roughly-isolated fibrinolytic component. It is approximately a 120 ml solution containing 300 mg enzyme.

5 (4) Put the above fibrinolytic component in a dialysis bag. Blow the bag with cold air till its volume is decreased to about 15 ml. Store at 4°C.

10 (5) Use Sephacryl S0200HR as gel chromatographic medium to pack the gel chromatographic column. The column volume is 2.7x70 cm³. Wash with 0.5 M NaCl solution for 1 hr at the flow rate of 99 ml/hr, then equilibrate with 0.15 M NaCl solution at the flow rate of 18 ml/min. The second elution peak is the refined antithrombosis enzyme solution which contains 170 mg enzyme in 70 ml of solution.

15 (6) Put the above refined antithrombosis enzyme into a dialysis bag. Dialyze for 3 hrs against distilled water and get 170 mg purified antithrombosis enzyme.

The purified enzyme was tested to have 1 fibrinolytic activity unit per mg protein. It was tested to show an

isoelectric point of pH 5.8 and MW of about 30 kD. No hemorrhagic side effect was detected with the enzyme.

Example 2

5 (1) Dissolve 4.5 g Southern-Anhui *Agkistrodon acutus* venom in 0.02 M, pH 7.8 phosphate buffer. The rest is the same as Step (1) of Example 1. Combine the supernatant and store at 4°C.

(2) Same as Step (2) of Example 1.

10 (3) Pack the column with the processed gel by the above step. Increase the column volume to 1600 ml (4x100 cm³). Keep the other conditions, 460 mg of enzyme was obtained after the chromatography.

15 (4) Put the roughly-isolated antithrombosis enzyme in a dialysis bag. Blow the bag with cold air till the volume is decreased to about 20 ml.

20 (5) Use Sephacryl S-300 HR as gel chromatographic medium to pack the chromatographic column. The column volume is 3.1x70 cm³. Wash 1 hr with 0.5 M NaCl at the flow rate of 99 ml/hr and equilibrate with 0.2 M NaCl. Load the concentrated

sample onto the column. Elute with 0.20 M NaCl solution at the flow rate of 24 ml/min. The second peak is the refined antithrombosis enzyme solution. Desalt the solution and get 280 mg purified enzyme in 100 ml solution.

5

Example 3

(1) Dissolve 3 g *Agkistrodon acutus* venom as described in Step (1) of Example 1.

(2) Use DEAE Sepharose Fast Flow as the anion exchanger. Process it and pack the column as described in Step (2) of Example 1. The column volume is 3x80 cm³. Rinse the column with 0.02 M pH 8.0 Tris-HCl buffer at the flow rate of 1.5 ml/min till equilibration. Load the sample and elute with a linear gradient. The elution buffer is the same as that in Step

(3) of Example 1. Adjust the flow rate to 80 ml/hr. 285 mg of enzyme was collected.

(3) The following procedure is the same as that of Example 1. In the end 160 mg refined antithrombosis enzyme in 60 ml solution was obtained which was shown to have fibrinolytic activity and no hemorrhagic side-effect.

Crystallization of the antithrombosis enzyme

The antithrombosis enzyme was obtained as described above (with a purity of 99% or higher) and further desalted. Crystals of the antithrombosis enzyme were grown by hanging-drop vapour diffusion technique using tissue culture plates and silicomised glass coverslips. At room temperature (about 20 °C), 3 µl of protein solution (30 mg/ml, in distilled water) was mixed with an equal volume of precipitant solution (0.1 M Mes buffer, pH 6.5, containing 16% PEG4000 (w/v) and 0.1 M CaAc₂). The solution was then equilibrated against 400 µl of precipitant solution containing 20% PEG 4000 (w/v). Crystals were collected from the solution in about one week. These crystals are useful for deciphering the tertiary structure and function center of the enzyme by X-ray crystallography and other techniques .

Characterization of the Antithrombosis Enzyme

The antithrombosis enzyme purified from the snake venom of Southern-Anhui *Agkistrodon acutus* contains Ca⁺⁺. Applicant studied the purity, fibrinolytic characteristic, antiplatelet

aggregation activity, amino acid structure, proteolytic activity and hemorrhagic activity of the antithrombosis enzyme.

5 The activities of the antithrombosis enzyme of the present invention may be assayed *in vitro* using any conventional technique. Preferably, the anti-thrombin assay involves direct determination of the thrombin-inhibitory activity of the molecule. Such techniques measure the inhibition of thrombin-catalyzed cleavage of colorimetric substrates or the increase in thrombin times or increase in activated partial thromboplastin times of human plasma. Alternatively, the assay employed may use purified thrombin and fibrinogen to measure the inhibition of release of fibrinopeptides.

15 The antiplatelet activity can be measured through a change in the degree of aggregation of platelets or a change in the release of a platelet secretory component in the presence of platelet activator. The former may be measured in an aggregometer. The latter may be measured using RIA or ELISA techniques specific for the secreted component.

Purity

Polyacrylamide gel electrophoresis of the enzyme as purified above under nonreducing condition showed one band at about 28-30 kD and the purity to be more than 90%.

Components

SDS-polyacrylamide gel electrophoresis was carried out according to the method of Laemmli (Nature 227:680-685,1970). Standards used for molecular weight determination were obtained from Sigma (bovine serum albumin - 66 kD, ovalbumin - 45 kD, glyceraldehyde-3-phosphate dehydrogenase - 36 kD, carbonic anhydrase - 29 kD, trypsinogen - 24 kD and lactalbumin - 14 kD). Staining was done with Coomassie brilliant blue R250.

Two bands were present on the SDS-PAGE gel. Thus the enzyme was isolated into two chains, A chain and B chain, each with a molecular weight of about 14 kD to about 16 kD on SDS-PAGE.

Fibrinolytic Activity

Fibrinolytic activity was measured with the fibrin-plate-clearance assay (Bajwa et al. Toxicon 18:285-290, 1980).

The positive control and the negative control were plasmin

(Sigma) and 0.9% sodium chloride, respectively. The experiment showed that the enzyme has high fibrinolytic activity in comparison to the control and the size of lysis area on the plate was proportional to the enzyme dosage. The enzyme was measured to hydrolyze fibrin at the level of about one to about three fibrinolytic activity units per mg of enzyme.

Fibrinogen cleavage sites

Fibrinogen was incubated with the antithrombosis enzyme to determine the initial cleavage sites. Fibrinogen (30 nmol) was dissolved in 2 ml of 0.05 M Tris buffer, pH 8.0. 10 μ l of the antithrombosis enzyme (0.6 nmol) was added to this solution. The solution was incubated at 37 °C for 1 hr. The reaction was stopped by the addition of 1 ml of 250 mM EDTA. This incubation time was employed as being optional for the accumulation of early degradation products.

The solution was then brought to 8 M Urea. Degraded fibrinogen fragments were reduced by the addition of dithiothreitol in a 10-fold molar excess over fibrinogen.

The sample was added to a C₁₈ reversed phase chromatography column for separation. Five fragments were

collected and freeze-dried. The amino-terminal amino acid sequences (1-10 residues) were determined. One fibrinogen cleavage site was found to be Met 235-Pro236 of the α chain of human fibrinogen. Other cleavage sites are still being studied.

5 Antiplatelet Aggregation Activity

ADP, collagen, epinephrine bitartrate and thrombin were obtained from Sigma Chemical Co. (St Louis, Mo). Venous blood was collected from healthy human donors who were drug-free and aspirin-free for at least 2 weeks before blood collection. Blood was collected into citrated Vacutainer tubes and centrifuged for 15 minutes at 150 g at room temperature. The platelet-rich plasma (PRP) was removed. The remaining blood was centrifuged for 15 minutes at 1500 g at room temperature, and the platelet-poor plasma (PPP) was removed.

15 Samples were assayed on a TYSM-91 Platelet Aggregometer with PPP as the blank (100% transmittance). PRP (200 μ l, 5×10^8 platelets/ml) was added to each micro test tube, and transmittance was set at 0%.

A stimulator of platelet aggregation such as ADP (200 μ l, 10 μ mol/l), collagen, epinephrine bitartrate or thrombin was added to a tube, and the aggregation profiles were plotted (percent transmittance versus time). The antithrombosis enzyme (20 μ l) of varying concentrations was added to the test tube before the addition of the platelet agonist. Antiplatelet aggregation activity is expressed as percent inhibition of agonist-induced platelet aggregation.

It was shown that the antithrombosis enzyme inhibits human platelet aggregation induced by ADP, Epinephrine, thrombin and collagen.

Primary Structure

The amino acid sequences of the A chain and the N-terminal of the B chain have been determined:

A:

Asp-Cys-Ser-Ser-Asp-Trp-Ser-Ser-Tyr-Glu-Gly-His-Cys-Tyr-Lys-Val-Phe-Lys-Gln-Ser-Lys-Thr-Trp-Thr-Asp-Ala-Glu-Ser-Phe-Cys-Thr-Lys-Gln-Val-Asn-Gly-Gly-His-Leu-Val-Ser-Ile-Glu-Ser-Ser-Gly-Glu-Ala-Asp-Phe-Val-Gly-Gln-Leu-Ile-Ala-Gln-Lys-Ile-Lys-Ser-Ala-Lys-Ile-His-Val-Trp-Ile-Gly-Leu-Arg-Ala-Gln-Asn-Lys-Glu-Lys-Gln-Cys-Ser-

Ile-Glu-Trp-Ser-Asp-Gly-Ser-Ser-Ile-Ser-Lys-Glu-Asn-Trp-Ile-Glu-
Glu-Glu-Ser-Lys-Lys-Cys-Leu-Gly-Val-His-Ile-Glu-Thr-Gly-Phe-His-
Lys-Trp-Glu-Asn-Phe-Tyr-Cys-Glu-Gln-Gln-Asp-Pro-Phe-Val-Cys-Glu-
Ala

5 B: N-

Asp-Cys-Pro-Ser-Glu-Trp-Ser-Ser-Tyr-Glu-Gly-Phe-Cys-Tyr-Lys-Pro-
Phe-

10 The amino acid sequence of the A chain was determined
by a combination of deduction from the cDNA sequence and amino
acid sequencing of the N-terminal of purified A chain.

15 Figure 1 shows the putative amino acid sequence of the
entire B chain, including a leader peptide, i.e., M G R F I
F V S F G L L V V F L S L S G T A A, which is
cleaved before the assembly of A chain and B chain to form the
antithrombosis enzyme.

Proteolytic Activity

20 The proteolytic activity of the antithrombosis enzyme
was assessed by azocasein hydrolysis. Azocasein was synthesized
from casein and diazotised sulfanilamide. The resulting
azocasein solution in 1% sodium bicarbonate had a concentration

of 42 mg/ml. 1 ml of the azocasein solution was added to a tube. A solution of the antithrombosis enzyme was added, and the tube was incubated at 37°C for 30 minutes. The perchloric acid was added into the tube, and the resulting precipitate was removed by centrifugation. Hydrolysis of azocasein was measured by increased absorbance at 390 nm of the supernatant. No proteolytic activity was detected in this assay.

Hemorrhagic Activity

The antithrombosis enzyme (100-200 µg) was injected subdermally under the clean shaven backs of white mice. After 24 hours, the mice were sacrificed and their skin observed for hemorrhage. No hemorrhage on the skin was observed after the administration of the antithrombosis enzyme.

Pharmacology of the Antithrombosis Enzyme in Animal Studies

Effect on Thrombosis

The antithrombosis enzyme was administered to rabbits intravenously. Thrombosis was determined before the administration and after the administration at 0.5, 1.0, 2.0, and 3.0 hours by the Chandler method. The enzyme showed anti-

thrombosis activity at 0.5 hr following administration at 0.005u/kg and this activity was increased significantly at 1.0 hr and at 0.01u/kg.

Effect on Arteria Thrombus and Venous Thrombus

5 Rats were anesthetized and their abdominal aorta and abdominal vein were exposed after surgery. Threads were passed through the abdominal aorta and abdominal vein. After two hours, the antithrombosis enzyme was injected from the femoral vein. Four hours later, the threads were taken out to observe the 10 arteria thrombus and the venous thrombus on the thread. The enzyme was shown to dissolve the arteria thrombus and the venous thrombus starting at 0.025u/kg and effectively at 0.05u/kg.

Effect on Pulmonary Thrombus

Pulmonary thrombus was detected and measured with ¹²⁵I 15 labeled fibrinogen. The radiation strength in the blood was determined with isotope counting method after the administration of the enzyme intravenously. The enzyme was shown to dissolve pulmonary thrombus in rats effectively at 0.005, 0.01, and 0.02 u/kg.

Effect on Fibrinogen

The enzyme was administered to rabbits intravenously. The content of fibrinogen in the plasma was determined at different times after administration. The enzyme was shown to
5 reduce the fibrinogen content in the blood at 0.0025u/kg, 0.005 u/kg, and 0.01 u/kg.

Effect on Platelet Aggregation

Dogs were anesthetized and placed on a respiratory pump with the stroke volume adjusted according to the dog's weight. The respirator dials were set to 20 beats per minute and 50%
10 respiration. Both femoral arteries were cannulated, one was attached to a transducer by which the heart beat and blood pressure were monitored on a chart recorder throughout the test, and the other was used for repeated bleeding sampling. The
15 formal vein was also cannulated to administer antithrombosis enzyme or vehicle and for the maintenance of anesthesia as needed.

Animals were dosed intravenously with antithrombosis enzyme at different dosage levels. Bleeding time and platelet
20 aggregation were determined before and at different intervals

after the administration of the antithrombosis enzyme. Aspirine (35 mg/kg) was used as the positive control and 0.9% sodium chloride was used as the negative control.

It was shown that the antithrombosis enzyme prolongs the bleeding time and inhibits platelet aggregation.

Effect on ELT

The enzyme was administered to rabbits intravenously. ELT in the blood of the rabbits was determined at different times after administration. The enzyme was shown to shorten ELT effectively starting at about 120 minutes from administration.

Effect on the Content of FDP in the Blood

The enzyme was administered to rabbits intravenously. The content of FDP in the blood of the rabbits was determined at different times after administration. The enzyme was shown to increase the content of FDP in the blood significantly in a dosage dependent manner.

Effect on The Whole Blood Clotting Time

The enzyme was administered to rabbits intravenously. The whole blood clotting time (CT) was determined. Two hours

following the administration, the enzyme showed anti-clotting activity.

Effect on the Kaolin Partial Thromboplastin Time

The enzyme was administered to rabbits intravenously.

5 The KPTT was determined. 240 minutes following the administration, the enzyme was shown to prolong the KPTT significantly.

Effect on Thrombin Time

The enzyme was administered to rabbits intravenously.

10 The TT was determined at the different time after administration. 240 minutes following the administration, the enzyme was shown to prolong the TT significantly.

Effect on Blood Viscosity

The enzyme was administered to rabbits intravenously.

15 The blood viscosity was determined at different times after administration. 120-240 minutes following the administration, the enzyme was shown to reduce the blood viscosity significantly.

General Pharmacology

20 Dogs were anesthetized with pentobarbital sodium (30 mg/kg). The antithrombosis enzyme at dosages in the range of

0.0015 - 0.06u/kg was administrated intravenously. The left
arteria carotis was isolated. The aortic systolic pressure,
diastolic pressure, mean pressure and heart rate were recorded
directly by TP-200T pressure transducer and RM-6000 multi-channel
5 physiological recorder. At the same time, the respiratory curve
and the electrocardiogram were recorded. The enzyme did not show
significant effects on the above parameters.

When the enzyme was administered to mice intravenously
from the tail, it did not affect the behavior of the mice. The
enzyme did not activate or restrain the neural system of the
mice.

Pharmacokinetics

The pharmacokinetics of the enzyme in the rats was
studied with ^{125}I labeled antithrombosis enzyme at 0.02, 0.01, and
15 0.005u/kg. The Concentration-Time Curves of the enzyme in the
blood were very similar at the three dosages which is in
agreement with the Two-Room Model. $T_{1/2}$ is 6 hours. Among
organs, the enzyme distributes mainly in the lung and the kidney.
The enzyme can infiltrate through the blood-brain-barrier. It is
20 excreted mainly through urine.

Cloning and Expression of the Antithrombosis Enzyme

A wide variety of methods may be used in locating and identifying cDNA sequences corresponding to A chain and B chain of this invention. The two most preferred techniques are the use of oligonucleotide probe based on the amino acid sequences of A chain and B chain and immunoscreening, which utilizes antibodies against A chain or B chain to detect clones which express cDNA sequences corresponding to the polypeptide.

The immuno screening technique requires that the cDNA library be contained in an expression vector. Such vectors include lambda gt11, lambda gt10 and other expression vectors known in the art. Antibodies employed in the immunoscreening technique include antibodies against the intact antithrombosis enzyme of the present invention, antibodies against denatured polypeptide and antibodies against peptide portions of the antithrombosis enzyme. Once a cDNA clone has been identified and isolated, it may be removed from the vector and analyzed to determine whether it contains the entire antithrombosis enzyme coding sequence. Partial cDNAs may themselves be used to reprobe the cDNA library and to locate full-length cDNAs.

The DNA molecules of this invention may be synthesized from oligonucleotides by chemical means using an oligonucleotide synthesizer. Such oligonucleotides may be designed based on the disclosed amino acid sequences of the antithrombosis enzyme.

5 Standard methods may be applied to synthesize a gene encoding the antithrombosis enzyme with knowledge of the enzyme's amino acid sequence. For example, the complete amino acid sequence may be used to construct a back-translated gene. A DNA oligomer containing a nucleotide sequence capable of coding for the desired antithrombosis enzyme may be synthesized in a single step. Alternatively, several smaller oligonucleotides coding for portions of the antithrombosis enzyme may be synthesized and subsequently ligated together. Preferably, the antithrombosis enzyme gene is synthesized as 10-20 separate oligonucleotides which are subsequently linked together. The individual
10
15 oligonucleotides contain 5' or 3' overhangs for complementary assembly.

Following synthesis of the oligomers and cleavage of the desired vector, assembly of the antithrombosis enzyme gene
20 may be achieved in one or more steps by techniques well known in

the art. Once assembled, the gene will be characterized by sequences which are recognized by restriction endonucleases, including unique restriction sites for direct assembly into a cloning or an expression vector; preferential codons based upon the host expression system to be used; and a sequence which, when transcribed, produces a mRNA with minimal secondary structure. Proper assembly maybe confirmed by nucleotide sequencing, restriction mapping, and expression of a biologically active antithrombosis enzyme in a suitable host.

It will be understood by those of skill in the art that, due to the degeneracy of the genetic code, many different synthetic DNAs will be capable of encoding the antithrombosis enzyme of this invention. It will also be apparent that many of these DNAs will be faithfully expressed in host transformed with them. Therefore, the present invention relates to not only one, but all DNA molecules which encode the desired antithrombosis enzyme and which can be expressed by one or more hosts transformed with them. Most of these DNA molecules will be capable of hybridizing to one another under moderately stringent conditions.

The DNA sequences and recombinant DNA molecules of the present invention may be inserted into and expressed using a wide variety of vectors. As such, the DNA sequence encoding the antithrombosis enzyme of the invention must be operatively linked to an expression control sequence. The term "operatively linked", as used herein refers to a positioning in a vector so that transcription and translation of the coding sequence is directed by the control sequence. Useful vectors may consist of segments of chromosomal, non-chromosomal and synthetic DNA sequences, such as various known derivatives of SV40, bovine papilloma virus, adenovirus and cytomegalovirus and known bacterial plasmids, e.g., plasmids from E. coli including colE1, pCR1, pBR322, pMB9 and RP4; phage DNAs, e.g., the numerous derivatives of lambda phage, e.g., NM 989, and other DNA phages, e.g., M13 and other Filamentous single-stranded DNA phages; vectors useful in yeasts, such as the 2 μ m plasmid; vectors useful in animal cells, such as those containing SV40 adenovirus and retrovirus-derived DNA sequences; commercially available high expression vectors, e.g., the pGEM series and the lambda Zap vectors; and vectors derived from combinations of plasmids and

phage DNAs, such as plasmids which have been modified to employ phage DNA or other derivatives thereof.

Such expression vectors are also characterized by at least one expression control sequence. When the DNA sequences of this invention are inserted in the vector they should be operatively linked to such expression control sequence in order to control and to regulate the expression of that cloned DNA sequence. Examples of useful expression control sequences include the malE system, the OmpA system, the lac system, the trp system, the tac system, the trc system, major operator and promoter regions of phage lambda, the control region of fd coat protein, the glycolytic promoters of yeast, e.g., the promoter for 3-phosphoglycerate kinase, the promoters of yeast acid phosphatase, e.g., Pho5, the promoters of the yeast-mating factors, and promoters derived from polyoma, adenovirus, retrovirus, and simian virus, e.g., the early and late promoters of SV40, and other sequences known to control the expression of genes of prokaryotic or eukaryotic cells and their viruses or combinations thereof.

Furthermore, within each specific expression vector, various sites may be selected for insertion of DNA sequences encoding the antithrombosis enzyme or a fusion protein of this invention. These sites are usually designated by the restriction
5 endonuclease which cuts them. They are well recognized by those of skill in the art. It is, of course, to be understood that an expression vector useful in this invention need not have a restriction endonuclease site for insertion of the chosen DNA fragment. Instead, the vector may be joined to the fragment by alternative means.

The expression vector, and in particular the site chosen therein for insertion of a selected DNA fragment and its operative linking therein to an expression control sequence, is determined by a variety of factors, e.g., number of sites
15 susceptible to a particular restriction enzyme, size of the protein to expressed, susceptibility of the desired protein to proteolytic degradation by host cell enzymes, contamination or binding of the protein to be expressed by host cell proteins difficult to remove during purification, expression
20 characteristics, such as the location of start and stop codons

relative to the vector sequences, and other factors recognized by those of skill in the art. The choice of a vector and an insertion site for a DNA sequence is determined by a balance of these factors, not all selections being equally effective for a given case.

Useful hosts which may be transformed with these vectors and which may be employed to express the polypeptides of this invention may include well known eukaryotic and prokaryotic hosts, such as strains of E. coli, such as E. coli SG-936, E. coli HB 101, E. coli W3110, E. coli X1776, E. coli X2282, E. coli DHI, and E. coli MRC1, Pseudomonas, Bacillus, such as bacillus subtilis, Streptomyces, yeasts and other fungi, animal cells, such as COS cells and CHO cells, human cells, insect cells and plant cells in tissue culture.

Of course, not all host/expression vector combinations will function with equal efficiency in expressing the DNA sequences of this invention or in producing the antithrombosis enzyme or fusion polypeptide. However, a particular selection of a host-expression vector combination may be made by those of skill in the art, after due consideration of the principles set

forth herein without departing from the scope of this invention. For example, the selection should be based on a balancing of a number of factors. These include, for example, compatibility of the host and vector, toxicity of the proteins encoded by the DNA
5 sequence to the host, ease of recovery of the desired protein, expression characteristics of the DNA sequences and the expression control sequences operatively linked to them, biosafety, costs and the folding, form or any other necessary post-expression modifications of the desired protein.

Constructing a λ cDNA library from the venom gland

After three days of venom extraction, *Agkistrodon* acutus snakes were sacrificed by decapitation. The venom gland was collected and immediately frozen in liquid nitrogen. Total RNA was extracted from the venom gland with isothiocyanate-
15 phenol-chloride. mRNA was purified from the total RNA by an oligo(dT)-cellulose column. The first DNA strand was synthesized by reverse transcriptase. The second DNA strand was then synthesized and EcoRI adapters were ligated to the ends of the cDNA. This cDNA was ligated into λ gt11 vector and packaged.
20 The packaged DNA was titered and found to contain 1.1×10^6

recombinants. Finally, 1.7×10^{10} pfu/ml were amplified to make the cDNA library.

Primers for cDNA amplification of the antithrombosis
polypeptides

5 One forward primer PF2 was designed according to the N-terminal amino acid sequence of the A chain.

PF2: Y E G H C Y, 5' TATGAAGGGCATTGCTACAA 3'

10 Another forward primer was designed from the 5'-terminal of the cDNA sequence of IX/X-Binding Protein containing the start codon (ATG):

PF3: 5' CCATGGGGCGATTCATCTTC 3'

15 A reverse primer (PR1) was designed from the 3'-terminal of the cDNA sequence of the IX/X-Binding Protein containing the stop codon (TGA).

PR1: 5' CAGCTGCATCTTCAGACTA 3'

20 The λ gt11 Forward Primer and Reverse Primer were also used to amplify the λ cDNA.

λ gt11 Forward Primer: 5' GGTGGCGACGACTCCTGGAGCCCCG 3'

λ gt11 Reverse Primer: 5' TTGACACCAGACCAACTGGTAATG 3'

Amplification of clones encoding the antithrombosis
polypeptides

The λ cDNA library was used as the template for PCR amplification. The PCR products were analyzed by 1.5% agarose gel electrophoresis and purified from low temperature melting gel. The PCR products were treated with Klenow fragment to generate blunt-ended DNA and inserted into the pBluscript vector at the SmaI site with the T4 DNA ligase. Fresh competent *E. coli* was transformed with the ligation solution and positive colonies were selected by color. The positive clones were digested with HindIII/BamHI and sequenced with T3 and T7 primers and the T7 sequencing Kit (Promega).

One of the clones amplified with PF2/poly(T) primers, Clone A, contains 544 bp as shown in SEQ ID NO: 1 which encodes the A chain of the antithrombosis enzyme from EGHCY to the carboxyl end.

One of the clone amplified with PF3/PR1 primers, Clone B, contains about 500 bp as shown in Figure 1 which putatively encodes the B chain of the antithrombosis enzyme.

The λ cDNA library was further screened with clone A as the probe. Several positive clones with inserts ranging from 500 bp to 1.2 kb have been selected to be sequenced.

5 Pharmaceutical Indications

Pathogenic platelet aggregation may yield thrombi (blood clots) which occlude blood flow to dependent tissues and lead to a variety of life-threatening vascular diseases, such as myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis, peripheral arterial occlusion and other blood system thromboses. Methods of preventing or inhibiting platelet aggregation and release are desirable in the treatment of these diseases. Inhibition of platelet aggregation may also be desirable in dialysis, storage of platelets in platelet concentrates, and following vascular surgery such as angioplasty.

The antithrombosis enzyme of this invention is effective for the treatment of thrombotic diseases, including, but not limited to, angiopathic thrombosis, cerebral thrombosis, acute thrombosis, peripheral angina pectoris, ischemic cerebral vascular disease, unstable angina, unstable stenocardia,

hemiparalysis caused by cerebral thrombosis, myocardial infarction, stroke, pulmonary embolism, deep vein thrombosis and peripheral arterial occlusion; restenosis following arterial injury or invasive cardiological procedures; acute or chronic atherosclerosis; edema and inflammation; abnormal cell regulatory processes (e.g. secretion, shape changes, proliferation); cancer and metastasis; and neurodegenerative diseases. It may also be used during the recovery period of cerebral hemorrhage to reduce or inhibit blood clot in the brain.

Toxicology Studies

Acute toxicity

The acute toxicity of the antithrombosis enzyme was studied with rats and mice and two injection routes (i.t.v. and i.p.). The results are shown below:

Table 1

animal	injection routes	LD ₅₀ (95% reliability)
mouse	i.t.v.	14.1u/kg (9.7~17.5u/kg)
	i.p.	23.3u/kg(18.4~29.5u/kg)
rat	i.t.v.	14.1u/kg(22.2~17.2u/kg)
	i.p.	14.1u/kg(46.7~63.5u/kg)

Long-term toxicity

For 8 weeks, rats were given the enzyme (i.t.v.) at dosages pharmaceutically equivalent to 12.5, 50, 200 times that's used clinically for adult humans. Neither abnormal behavior nor toxic symptom was observed. The consumption of food and water, net weight increase, various biochemical indexes and coefficients of major organs and tissues showed no visible change among the testing groups. Pathological histology examination of major organs did not show distinct toxicosis pathology changes.

Side Effects & Contra-indications:

The enzyme has no neural toxicity and is non-hemorrhagic. No obvious side effects have been noticed in clinical use.

The antithrombosis enzyme polypeptides, antibodies and hybridomas

A variety of methodologies known in the art can be utilized to obtain the polypeptide of the present invention. The polypeptide may be purified from a snake which naturally produces the polypeptide, chemically synthesized, or expressed by recombinant techniques.

Any eukaryotic organism can be used as a source for the polypeptide of the invention, as long as the source organism expresses such a polypeptide. One skilled in the art can readily follow known methods for isolating proteins in order to obtain the polypeptide free of natural contaminants. These include, but are not limited to: size-exclusion chromatography, HPLC, ion-exchange chromatography, and immuno-affinity chromatography.

The present invention relates to an antibody having binding affinity to the antithrombosis enzyme polypeptide. The present invention also relates to an antibody having specific binding affinity to the antithrombosis enzyme polypeptide. Such an antibody may be isolated by comparing its binding affinity to the antithrombosis enzyme polypeptide with its binding affinity to another polypeptide. Those which bind selectively to the antithrombosis enzyme would be chosen for use in methods requiring a distinction between the antithrombosis enzyme and other polypeptides.

The antithrombosis enzyme polypeptide of the present invention can be used in a variety of procedures and methods, such as for the generation of antibodies, for use in identifying

pharmaceutical compositions, and for studying DNA/protein interaction.

The antithrombosis enzyme polypeptide of the present invention can be used to produce antibodies or hybridomas. One skilled in the art will recognize that if an antibody is desired, such a peptide would be generated as described herein and used as an immunogen. The antibodies of the present invention include monoclonal and polyclonal antibodies, as well fragments of these antibodies, and humanized forms. Humanized forms of the antibodies of the present invention may be generated using one of the procedures known in the art such as chimerization or CDR grafting. The present invention also relates to a hybridoma which produces the above-described monoclonal antibody, or binding fragment thereof. A hybridoma is an immortalized cell line which is capable of secreting a specific monoclonal antibody.

In general, techniques for preparing monoclonal antibodies and hybridomas are well known in the art (Campbell, Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers,

Amsterdam, The Netherlands (1984); St. Groth et al., J. Immunol. Methods 35:1-21(1980)). Any animal (mouse, rabbit, and the like) which is known to produce antibodies can be immunized with the selected polypeptide. Methods for immunization are well known in the art. Such methods include subcutaneous or intraperitoneal injection of the polypeptide. One skilled in the art will recognize that the amount of polypeptide used for immunization will vary based on the animal which is immunized, the antigenicity of the polypeptide and the site of injection.

The polypeptide may be modified or administered in an adjuvant in order to increase the peptide antigenicity. Methods of increasing the antigenicity of a polypeptide are well known in the art. Such procedures include coupling the antigen with a heterologous protein (such as globulin or β -galactosidase) or through the inclusion of an adjuvant during immunization.

For monoclonal antibodies, spleen cells from the immunized animals are removed, fused with myeloma cells, such as SP2/0-Agl4 myeloma cells, and allowed to become monoclonal antibody producing hybridoma cells. Any one of a number of methods well known in the art can be used to identify the

hybridoma cell which produces an antibody with the desired characteristics. These include screening the hybridomas with an ELISA assay, western blot analysis, or radioimmunoassay (Lutz et al., Exp. Cell Res. 175:109-124(1988)). Hybridomas secreting
5 the desired antibodies are cloned and the class and subclass is determined using procedures known in the art (Campbell, Monoclonal Antibody Technology: Laboratory Techniques in Biochemistry and Molecular Biology, supra (1984)).

For polyclonal antibodies, antibody containing antisera
10 are isolated from the immunized animal and screened for the presence of antibodies with the desired specificity using one of the above-described procedures. The above-described antibodies may be detectably labeled. Antibodies can be detectably labeled through the use of radioisotopes, affinity labels (such as
15 biotin, avidin, and the like), enzymatic labels (such as horse radish peroxidase, alkaline phosphatase, and the like) fluorescent labels (such as FITC or rhodamine, and the like), paramagnetic atoms, and the like. Procedures for accomplishing such labeling are well-known in the art, for example, see
20 (Stemberger et al., J. Histochem. Cytochem. 18:315(1970);

5 Bayer et al., Meth. Enzym. 62:308(1979); Engval et al.,
Immunot. 109:129(1972); Goding, J. Immunol. Meth.
13:215(1976)). The labeled antibodies of the present invention
can be used for *in vitro*, *in vivo*, and *in situ* assays to identify
cells or tissues which express a specific peptide.

10 The above-described antibodies may also be immobilized
on a solid support. Examples of such solid supports include
plastics such as polycarbonate, complex carbohydrates such as
agarose and sepharose, acrylic resins and such as polyacrylamide
and latex beads. Techniques for coupling antibodies to such
solid supports are well known in the art (Weir et al., Handbook
of Experimental Immunology 4th Ed., Blackwell Scientific
Publications, Oxford, England, Chapter 10(1986); Jacoby et al.,
15 Meth. Enzym. 34 Academic Press, N.Y. (1974)). The immobilized
antibodies of the present invention can be used for *in vitro*, *in*
vivo, and *in situ* assays as well as in immunochromotography.

20 Furthermore, one skilled in the art can readily adapt
currently available procedures, as well as the techniques,
methods and kits disclosed above with regard to antibodies, to
generate peptides capable of binding to a specific peptide

sequence in order to generate rationally designed antipeptide
peptides, for example see Hurby et al., Application of Synthetic
Peptides: Antisense Peptides, In Synthetic Peptides, A User's
Guide, W.H. Freeman, NY, pp. 289-307(1992), and Kaspczak et
5 al., Biochemistry 28:9230-8(1989).

The present invention encompasses a method of detecting
the antithrombosis enzyme polypeptide in a sample, comprising: a)
contacting the sample with an above-described antibody, under
conditions such that immunocomplexes form, and b) detecting the
10 presence of said antibody bound to the polypeptide. In detail,
the methods comprise incubating a test sample with one or more of
the antibodies of the present invention and assaying whether the
antibody binds to the test sample.

Conditions for incubating an antibody with a test
15 sample vary. Incubation conditions depend on the format employed
in the assay, the detection methods employed, and the type and
nature of the antibody used in the assay. One skilled in the art
will recognize that any one of the commonly available
immunological assay formats (such as radioimmunoassays, enzyme-
20 linked immunosorbent assays, diffusion based Ouchterlony, or

rocket immunofluorescent assays) can readily be adapted to employ the antibodies of the present invention. Examples of such assays can be found in Chard, An Introduction to Radioimmunoassay and Related Techniques Elsevier Science Publishers, Amsterdam, The Netherlands (1986); Bullock et al., Techniques in Immunocytochemistry, Academic Press, Orlando, FL Vol. 1(1982), Vol. 2 (1983), Vol. 3 (1985); Tijssen, Practice and Theory of Enzyme Immunoassays: Laboratory Techniques in Biochemistry and Molecular Biology, Elsevier Science Publishers, Amsterdam, The Netherlands (1985).

The immunological assay test samples of the present invention include cells, protein or membrane extracts of cells, or biological fluids such as blood, serum, plasma, or urine. The test sample used in the above-described method will vary based on the assay format, nature of the detection method and the tissues, cells or extracts used as the sample to be assayed. Methods for preparing protein extracts or membrane extracts of cells are well known in the art and can be readily be adapted in order to obtain a sample which is capable with the system utilized.

This invention relates to a kit containing all the necessary reagents to carry out the previously described methods of detection. The kit may comprise: i) a first container means containing an above-described antibody, and ii) second container means containing a conjugate comprising a binding partner of the antibody and a label. In another preferred embodiment, the kit further comprises one or more other containers comprising one or more of the following: wash reagents and reagents capable of detecting the presence of bound antibodies.

Examples of detection reagents include, but are not limited to, labeled secondary antibodies, or in the alternative, if the primary antibody is labeled, the chromophoric, enzymatic, or antibody binding reagents which are capable of reacting with the labeled antibody. The compartmentalized kit may be as described above for nucleic acid probe kits. One skilled in the art will readily recognize that the antibodies described in the present invention can readily be incorporated into one of the established kit formats which are well known in the art.

Pharmaceutical formulations and modes of administration

The antithrombosis enzyme of the present invention may be formulated into pharmaceutically acceptable compositions for inhibiting both thrombin- and platelet mediated functions in a patient or in extracorporeal blood.

The dosage and dose rate of the antithrombosis enzyme of this invention will depend on a variety of factors, such as the patient's body weight, the specific pharmaceutical composition used, and the object of the treatment, i.e., therapy or prophylaxis, the nature of the thrombotic disease to be treated, and the judgment of the treating physician. A pharmaceutically effective amount of the antithrombosis enzyme of this invention will normally be in the dosage range of between about 0.001-500 mg/kg body weight, preferably about 0.1-50 mg/kg body weight. For the treatment of extracorporeal blood, the antithrombosis enzyme should be used at about 0.005-50 μ g/ml, preferably at about 0.5-5 μ g/ml of extracorporeal blood. It should be understood that other dosages outside of these illustrative ranges may be employed in the pharmaceutical compositions of this invention.

The antithrombosis enzyme that affects the disorders of interest can be administered in pharmaceutically acceptable formulations where it is mixed with suitable carriers or excipient(s). These formulations can be administered by standard routes, including, but not limited to, intracerebroventricular, intracerebral, intravaginal, intrauterine or parenteral (e.g., intravenous, intraspinal, subcutaneous or intramuscular) route.

In treating a patient exhibiting a disorder of interest, a therapeutically effective amount of a agent or agents is administered. A therapeutically effective dose refers to that amount of the compound that results in amelioration of symptoms or a prolongation of survival in a patient.

Toxicity and therapeutic efficacy of such compounds can be determined by standard pharmaceutical procedures in cell cultures or experimental animals, e.g., for determining the LD₅₀ (the dose lethal to 50% of the population) and the ED₅₀ (the dose therapeutically effective in 50% of the population). The dose ratio between toxic and therapeutic effects is the therapeutic index and it can be expressed as the ratio LD₅₀/ED₅₀. Compounds which exhibit large therapeutic indices are preferred. The data

obtained from these cell culture assays and animal studies can be used in formulating a range of dosage for use in human. The dosage of such compounds lies preferably within a range of circulating concentrations that include the ED_{50} with little or no toxicity. The dosage may vary within this range depending upon the dosage form employed and the route of administration utilized.

For any compound used in the method of the invention, the therapeutically effective dose can be estimated initially from cell culture assays. For example, a dose can be formulated in animal models to achieve a circulating plasma concentration range that includes the IC_{50} as determined in cell culture (i.e., the concentration of the test compound which achieves a half-maximal disruption of the protein complex, or a half-maximal inhibition of the cellular level and/or activity of a complex component). Such information can be used to more accurately determine useful doses in humans. Levels in plasma may be measured, for example, by HPLC.

The dosage of the antithrombosis enzyme will depend on the disease state or condition being treated and other clinical

factors such as weight and condition of the patient and the route of administration of the compound. It is to be understood that the present invention has application for both human and veterinary use. The present invention contemplates single as well as multiple administrations, given either simultaneously or over an extended period of time.

The exact formulation, route of administration and dosage can be chosen by the individual physician in view of the patient's condition. (See e.g. Fingl et al., in The Pharmacological Basis of Therapeutics, 1975, Ch. 1 p. 1). It should be noted that the attending physician would know how to and when to terminate, interrupt, or adjust administration due to toxicity, or to organ dysfunctions. Conversely, the attending physician would also know to adjust treatment to higher levels if the clinical response were not adequate (precluding toxicity). The magnitude of an administered dose in the management of the oncogenic disorder of interest will vary with the severity of the condition to be treated and to the route of administration. The severity of the condition may, for example, be evaluated, in part, by standard prognostic evaluation methods. Further, the

dose and perhaps dose frequency, will also vary according to the age, body weight, and response of the individual patient. A program comparable to that discussed above may be used in veterinary medicine.

5 Depending on the specific conditions being treated, such agents may be formulated and administered systemically or locally. Techniques for formulation and administration may be found in Remington's Pharmaceutical Sciences, 18th ed., Mack Publishing Co., Easton, PA (1990). Suitable routes may include
10 intestinal administration or parenteral delivery, including intramuscular, subcutaneous, intramedullary injections, as well as intrathecal, direct intraventricular, intravenous, intraperitoneal, intranasal, or intraocular injections, just to name a few.

15 For injection, the agents of the invention may be formulated in aqueous solutions, preferably in physiologically compatible buffers such as Hanks's solution, Ringer's solution, or physiological saline buffer. For such transmucosal administration, penetrants appropriate to the barrier to be

permeated are used in the formulation. Such penetrants are generally known in the art.

Use of pharmaceutically acceptable carriers to formulate the compounds herein disclosed for the practice of the invention into dosages suitable for systemic administration is within the scope of the invention. With proper choice of carrier and suitable manufacturing practice, the compositions of the present invention, in particular, those formulated as solutions, may be administered parenterally, such as by intravenous injection. The compounds can be formulated readily using pharmaceutically acceptable carriers well known in the art into dosages suitable for oral administration. Such carriers enable the compounds of the invention to be formulated as tablets, pills, capsules, liquids, gels, syrups, slurries, suspensions and the like, for oral ingestion by a patient to be treated.

Agents intended to be administered intracellularly may be administered using techniques well known to those of ordinary skill in the art. For example, such agents may be encapsulated into liposomes, then administered as described above. Liposomes are spherical lipid bilayers with aqueous interiors. All

molecules present in an aqueous solution at the time of liposome formation are incorporated into the aqueous interior. The liposomal contents are both protected from the external microenvironment and, because liposomes fuse with cell membranes, are efficiently delivered into the cell cytoplasm. Additionally, due to their hydrophobicity, small organic molecules may be directly administered intracellularly.

Pharmaceutical compositions suitable for use in the present invention include compositions wherein the active ingredients are contained in an effective amount to achieve its intended purpose. Determination of the effective amounts is well within the capability of those skilled in the art, especially in light of the detailed disclosure provided herein. In addition to the active ingredients, these pharmaceutical compositions may contain suitable pharmaceutically acceptable carriers comprising excipients and auxiliaries which facilitate processing of the active compounds into preparations which can be used pharmaceutically. The preparations formulated for oral administration may be in the form of tablets, dragees, capsules, or solutions. The pharmaceutical compositions of the present invention may be

manufactured in a manner that is itself known, e.g., by means of conventional mixing, dissolving, granulating, dragee-making, levitating, emulsifying, encapsulating, entrapping or lyophilizing processes.

5 Pharmaceutical formulations for parenteral administration include aqueous solutions of the active compounds in water-soluble form. Additionally, suspensions of the active compounds may be prepared as appropriate oily injection suspensions. Suitable lipophilic solvents or vehicles include
10 fatty oils such as sesame oil, or synthetic fatty acid esters, such as ethyl oleate or triglycerides, or liposomes. Aqueous injection suspensions may contain substances which increase the viscosity of the suspension, such as sodium carboxymethyl cellulose, sorbitol, or dextran. Optionally, the suspension may
15 also contain suitable stabilizers or agents which increase the solubility of the compounds to allow for the preparation of highly concentrated solutions.

Dosage & Parental Administration

20 The antithrombosis enzyme could be administered to patients by intravenous injection. The enzyme is supplied as

sterile lyophilized white powder in vials. Each vial contains 0.25 unit. For treatment, dissolve this drug aseptically by adding proper sterile water. The solution is diluted with 250 ml of 0.9% Sodium Chloride and is given intravenously at the rate of 45 drops/min. 0.25 unit is given per day.

For acute treatment 0.5u is given on the first day and 0.25u/day afterwards. The course of treatment is two weeks.

The drug should be administered under the supervision of a physician.

Perform intradermal test before administration. The drug is diluted into 0.0025u/ml. Take 0.1 ml and inject intradermally. Observe the result after 20 min. The drug should not be given to the patient if the test result is positive.

Platelet count should be checked in regular intervals (once a week). If the platelet count falls below 80,000/mm³, the treatment should be discontinued immediately.

Gene Therapy with the ATE

The antithrombosis enzyme or its genetic sequences will be useful in gene therapy (reviewed in Miller, Nature 357:455-

460, 1992). Miller states that advances have resulted in practical approaches to human gene therapy that have demonstrated positive initial results. An *in vivo* model of gene therapy for human severe combined immunodeficiency is described in Ferrari, et al., *Science* 251:1363-1366, (1991). The basic science of gene therapy is described in Mulligan, *Science* 260:926-931, (1993).

In one preferred embodiment, one or more expression vector containing the ATE coding sequence is inserted into cells, the cells are grown *in vitro* and then infused in large numbers into patients.

The gene therapy may involve the use of an adenovirus containing ATE cDNA targeted to a vascular tissue, systemic ATE increase by implantation of engineered cells, or injection of naked ATE DNA into appropriate tissues.

Expression vectors derived from viruses such as retroviruses, vaccinia virus, adenovirus, adeno-associated virus, herpes viruses, several RNA viruses, or bovine papilloma virus, may be used for delivery of nucleotide sequences (e.g., cDNA) encoding recombinant ATE polypeptides into the targeted cell population (e.g., tumor cells). Methods which are well known to

those skilled in the art can be used to construct recombinant viral vectors containing coding sequences. See, for example, the techniques described in Maniatis et al., Molecular Cloning: A Laboratory Manual, Cold Spring Harbor Laboratory, N.Y. (1989), and in Ausubel et al., Current Protocols in Molecular Biology, Greene Publishing Associates and Wiley Interscience, N.Y. (1989). Alternatively, recombinant nucleic acid molecules encoding protein sequences can be used as naked DNA or in reconstituted system e.g., liposomes or other lipid systems for delivery to target cells (See e.g., Felgner et al., *Nature* 337:387-8, 1989). Several other methods for the direct transfer of plasmid DNA into cells exist for use in human gene therapy and involve targeting the DNA to receptors on cells by complexing the plasmid DNA to proteins. See, Miller, *supra*.

In its simplest form, gene transfer can be performed by simply injecting minute amounts of DNA into the nucleus of a cell, through a process of microinjection. Capecchi MR, Cell 22:479-88 (1980). Once recombinant genes are introduced into a cell, they can be recognized by the cell's normal mechanisms for transcription and translation, and a gene product will be

expressed. Other methods have also been attempted for introducing DNA into larger numbers of cells. These methods include: transfection, wherein DNA is precipitated with CaPO_4 and taken into cells by pinocytosis (Chen C. and Okayama H, Mol. Cell Biol. 7:2745-52 (1987)); electroporation, wherein cells are exposed to large voltage pulses to introduce holes into the membrane (Chu G. et al., Nucleic Acids Res., 15:1311-26 (1987)); lipofection/liposome fusion, wherein DNA is packaged into lipophilic vesicles which fuse with a target cell (Felgner PL., et al., Proc. Natl. Acad. Sci. USA. 84:7413-7 (1987)); and particle bombardment using DNA bound to small projectiles (Yang NS. et al., Proc. Natl. Acad. Sci. 87:9568-72 (1990)). Another method for introducing DNA into cells is to couple the DNA to chemically modified proteins.

It has also been shown that adenovirus proteins are capable of destabilizing endosomes and enhancing the uptake of DNA into cells. The admixture of adenovirus to solutions containing DNA complexes, or the binding of DNA to polylysine covalently attached to adenovirus using protein crosslinking agents substantially improves the uptake and expression of the

recombinant gene. Curiel DT et al., Am. J. Respir. Cell. Mol. Biol., 6:247-52 (1992).

As used herein "gene transfer" means the process of introducing a foreign nucleic acid molecule into a cell. Gene transfer is commonly performed to enable the expression of a particular product encoded by the gene. The product may include a protein, polypeptide, anti-sense DNA or RNA, or enzymatically active RNA. Gene transfer can be performed in cultured cells or by direct administration into animals. Generally gene transfer involves the process of nucleic acid contact with a target cell by non-specific or receptor mediated interactions, uptake of nucleic acid into the cell through the membrane or by endocytosis, and release of nucleic acid into the cytoplasm from the plasma membrane or endosome. Expression may require, in addition, movement of the nucleic acid into the nucleus of the cell and binding to appropriate nuclear factors for transcription.

As used herein "gene therapy" is a form of gene transfer and is included within the definition of gene transfer as used herein and specifically refers to gene transfer to

express a therapeutic product from a cell *in vivo* or *in vitro*.

Gene transfer can be performed *ex vivo* on cells which are then transplanted into a patient, or can be performed by direct administration of the nucleic acid or nucleic acid-protein complex into the patient.

In another preferred embodiment, a vector having nucleic acid sequences encoding ATE is provided in which the nucleic acid sequence is expressed only in specific tissue. Methods of achieving tissue-specific gene expression as set forth in International Publication No. WO 93/09236, filed November 3, 1992 and published May 13, 1993.

In all of the preceding vectors set forth above, a further aspect of the invention is that the nucleic acid sequence contained in the vector may include additions, deletions or modifications to some or all of the sequence of the nucleic acid, as defined above.

In another preferred embodiment, a method of gene replacement is set forth. "Gene replacement" as used herein means supplying a nucleic acid sequence which is capable of being expressed *in vivo* in an animal and thereby providing or

5

5

Other embodiments of this invention are disclosed in the following claims.

SEQUENCE LISTING

(1) GENERAL INFORMATION:

(i) APPLICANT: Chun Wang, Benjamin XY Li, Xin Cheng,
Jing Liu, Li-Wen Niu, Wan-Zhi Huang,
Zhen-Yu Xu, Dan Luo, Lian-Di Kang,
Jin-Guo Ding, Fang Rong, Yan Liu,
Hui-Ran Chen

(ii) TITLE OF INVENTION: AN ANTITHROMBOSIS ENZYME FROM THE
SNAKE VENOM OF AGKISTRODON ACUTUS

(iii) NUMBER OF SEQUENCES: 4

(iv) CORRESPONDENCE ADDRESS:

(A) ADDRESSEE: Lyon & Lyon
(B) STREET: 633 West Fifth Street
(C) CITY: Los Angeles
(D) STATE: California
(E) COUNTRY: USA
(F) ZIP: 90071

(v) COMPUTER READABLE FORM:

(A) MEDIUM TYPE: 3.5" Diskette, 1.44 Mb
(B) COMPUTER: IBM compatible
(C) OPERATING SYSTEM: IBM P.C. DOS (Version 5.0)
(D) SOFTWARE: WordPerfect (Version 5.1)

(vi) CURRENT APPLICATION DATA:

(A) APPLICATION NUMBER:
(B) FILING DATE:
(C) CLASSIFICATION:

(viii) ATTORNEY/AGENT INFORMATION:

(A) NAME: Chen, Anthony C.
(B) REGISTRATION NUMBER: 38,673

(C) REFERENCE/DOCKET NUMBER: 233/298

(ix) TELECOMMUNICATION INFORMATION:

5 (A) TELEPHONE: (213) 489-1600
(B) TELEFAX: (213) 955-0440
(C) TELEX: 67-3510

(2) INFORMATION FOR SEQ ID NO: 1:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 544
(B) TYPE: nucleic acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(xi) SEQUENCE DESCRIPTION : SEQ ID NO: 1:

GAAGGGCATT GCTACAAGGT CTTCAAACAA TCTAAGACCT GGACAGATGC 50
AGAGAGCTTC TGCACGAAGC AGGTGAACGG GGGGCATCTG GTCTCTATCG 100
AAAGCTCCGG AGAAGCAGAC TTTGTGGGCC AGTTGATTGC TCAGAAGATA 150
AAGTCAGCCA AAATCCATGT CTGGATCGGA CTGAGGGCTC AAAACAAAGA 200
AAAGCAATGC AGCATAGAGT GGAGCGATGG CTCCAGCATC AGTTATGAGA 250
ATTGGATTGA AGAAGAATCC AAAAAGTGTC TTGGGGTGCA CATAGAGACA 300
GGGTTTCATA AGTGGGAGAA TTTTACTGT GAACAACAAG ATCCTTTTGT 350
CTGCGAGGCA TAGTCTGAAG ATCCAGCTGA TTGAAGTCTG GAGAAGCAAG 400
GAAGCCCCCC ACCCCATCCC CCAACCCTGC CTAGCCACAA TCTCTGCTAT 450
GCACCCTTTG CTCAACGGAT GCTCTCTGTA GCTGGATCTG GTGTTGCTGC 500
TCCTGATGGG CCGGAAGTCA ATAAATTCTG CCTAGCCTGA AAAA 544

(2) INFORMATION FOR SEQ ID NO: 2:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 129

(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

5 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION : SEQ ID NO: 2

10 Asp Cys Ser Ser Asp Trp Ser Ser Tyr Glu Gly His Cys Tyr Lys
Val Phe Lys Gln Ser Lys Thr Trp Thr Asp Ala Glu Ser Phe Cys
Thr Lys Gln Val Asn Gly Gly His Leu Val Ser Ile Glu Ser Ser
15 Gly Glu Ala Asp Phe Val Gly Gln Leu Ile Ala Gln Lys Ile Lys
Ser Ala Lys Ile His Val Trp Ile Gly Leu Arg Ala Gln Asn Lys
Glu Lys Gln Cys Ser Ile Glu Trp Ser Asp Gly Ser Ser Ile Ser
20 Lys Glu Asn Trp Ile Glu Glu Glu Ser Lys Lys Cys Leu Gly Val
His Ile Glu Thr Gly Phe His Lys Trp Glu Asn Phe Tyr Cys Glu
25 Gln Gln Asp Pro Phe Val Cys Glu Ala

(2) INFORMATION FOR SEQ ID NO: 3:

30 (i) SEQUENCE CHARACTERISTICS:
(A) LENGTH: 29
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

35 (ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION : SEQ ID NO: 3

40 Asp Cys Ser Ser Asp Trp Ser Ser Tyr Glu Gly His Cys Tyr Lys

Val Phe Lys Gln Ser Lys Thr Trp Thr Asp Ala Glu Ser Phe

(2) INFORMATION FOR SEQ ID NO: 4:

(i) SEQUENCE CHARACTERISTICS:

(A) LENGTH: 17
(B) TYPE: amino acid
(C) STRANDEDNESS: single
(D) TOPOLOGY: linear

(ii) MOLECULE TYPE: peptide

(xi) SEQUENCE DESCRIPTION : SEQ ID NO: 4

Asp Cys Pro Ser Glu Trp Ser Ser Tyr Glu Gly Phe Cys Tyr Lys

Pro Phe